

IDENTIFICATION THROUGH INDUCTIVE VERIFICATION APPLICATION TO MONOTONE QUANTIFIERS

Nina Gierasimczuk

Institute for Logic, Language and Computation
Universiteit van Amsterdam

Logic and Intelligent Interaction
ESSLLI08
August 11, 2008

OUTLINE

- 1 PROBLEMS
- 2 QUANTIFIERS
 - Quantifiers of type $\langle 1 \rangle$
 - Quantifiers of type $\langle 1, 1 \rangle$
- 3 COMPUTATIONAL EPISTEMOLOGY
- 4 IDENTIFIABILITY
- 5 GENERAL QUESTION



PLAN

1 PROBLEMS

2 QUANTIFIERS

- Quantifiers of type $\langle 1 \rangle$
- Quantifiers of type $\langle 1, 1 \rangle$

3 COMPUTATIONAL EPISTEMOLOGY

4 IDENTIFIABILITY

5 GENERAL QUESTION

PROBLEMS

- Epistemological properties of quantifiers.
- Their influence on NL comprehension.
- Linking them to learnability features.
- Compare notions of decidability and identifiability.

PLAN

1 PROBLEMS

2 QUANTIFIERS

- Quantifiers of type $\langle 1 \rangle$
- Quantifiers of type $\langle 1, 1 \rangle$

3 COMPUTATIONAL EPISTEMOLOGY

4 IDENTIFIABILITY

5 GENERAL QUESTION



QUANTIFIERS

LINDSTRÖM DEFINITION

DEFINITION

A monadic generalized quantifier of type $\underbrace{\langle 1, \dots, 1 \rangle}_n$ is a class Q of structures of the form $\mathbf{M} = (M, A_1, \dots, A_n)$, where A_i is a subset of M . Additionally, Q is closed under isomorphism.



Q OF TYPE $\langle 1 \rangle$

MONOTONICITY

DEFINITION

Q_M is $\text{MON}\uparrow$ iff: if $A \subseteq A' \subseteq M$, then $Q_M(A)$ implies $Q_M(A')$.

DEFINITION

Q_M is $\text{MON}\downarrow$ iff: if $A' \subseteq A \subseteq M$, then $Q_M(A)$ implies $Q_M(A')$.



Q OF TYPE $\langle 1 \rangle$

EXTENDABILITY

DEFINITION

Q of type $\langle 1 \rangle$ satisfies *EXT* iff for all models \mathbf{M} and \mathbf{M}' :
 $A \subseteq M \subseteq M'$ implies $Q_{\mathbf{M}}(A) \iff Q_{\mathbf{M}'}(A)$.



QUANTIFIERS OF TYPE $\langle 1, 1 \rangle$

PRECONDITIONS

Restriction to CE quantifiers.

DEFINITION

Let Q be of type $\langle 1, 1 \rangle$.

Then for all \mathbf{M}, \mathbf{M}' , all $A, B \subseteq M$, and $A', B' \subseteq M'$:

(ISOM) If $(M, A, B) \cong (M', A', B')$, then $Q_{\mathbf{M}}(A, B) \Leftrightarrow Q_{\mathbf{M}'}(A', B')$.


(CONS) $Q_{\mathbf{M}}(A, B) \Leftrightarrow Q_{\mathbf{M}}(A, A \cap B)$.

(EXT) If $M \subseteq M'$, then $Q_{\mathbf{M}}(A, B) \Leftrightarrow Q_{\mathbf{M}'}(A, B)$.




CE QUANTIFIERS

(ISOM) IF $(M, A, B) \cong (M', A', B')$, THEN $Q_M(A, B) \Leftrightarrow Q_{M'}(A', B')$




CE QUANTIFIERS - EXT

(EXT) IF $M \subseteq M'$, THEN $Q_M(A, B) \Leftrightarrow Q_{M'}(A, B)$





CE QUANTIFIERS - CONS

(CONS) $Q_M(A, B) \Leftrightarrow Q_M(A, A \cap B)$



CE QUANTIFIERS

The scope we are interested in for both $\langle 1 \rangle$ and $\langle 1, 1 \rangle$ cases.



Q OF TYPE $\langle 1, 1 \rangle$

MONOTONICITY

DEFINITION

Q of type $\langle 1, 1 \rangle$ is $\text{MON}\uparrow$ iff:

If $A \subseteq M$ and $B \subseteq B' \subseteq M$, then $Q_M(A, B) \Rightarrow Q_M(A, B')$.



Q OF TYPE $\langle 1, 1 \rangle$

PERSISTENCE

DEFINITION

Q of type $\langle 1, 1 \rangle$ is PER iff:

If $A \subseteq A' \subseteq M$ and $B \subseteq M$, then $Q_M(A, B) \Rightarrow Q_M(A', B)$.



EXAMPLES

Determiner	MON \uparrow	EXT (for $\langle 1 \rangle$)	PER (for $\langle 1, 1 \rangle$)
All	+	-	-
No	-	-	-
Some	+	+	+
At least n	+	+	+
At most n	-	+	-
Exactly n	-	+	-

TABLE: Quantifiers and their properties



MONOTONICITY & LINGUISTICS

- Does monotonicity influence NL comprehension?
- Does monotonicity influence NL learning?
- Monotonicity and inference patterns (B. Geurts).
- Proposal: focus on persistence.

PLAN

- 1 PROBLEMS
- 2 QUANTIFIERS
 - Quantifiers of type $\langle 1 \rangle$
 - Quantifiers of type $\langle 1, 1 \rangle$
- 3 COMPUTATIONAL EPISTEMOLOGY**
- 4 IDENTIFIABILITY
- 5 GENERAL QUESTION

LOGIC OF RELIABLE INQUIRY - KEVIN KELLY

- Inspired by learning theory.
- Similar framework.
- Verification/falsification in computational setting.

LOGIC OF RELIABLE INQUIRY - KEVIN KELLY

ε — infinite string of data;

$\varepsilon|n$ — finite initial segment of ε through the position $n - 1$;

h — hypothesis;

C — correctness relation;

$C(\varepsilon, h)$ iff h is correct w.r.t. ε ;

α — an assessment method;

OUT conjectures 1, 0, !.



CERTAINTY IN RELIABLE INQUIRY

DEFINITION

α produces b with certainty on (h, ε) iff there is an n s.t.:

- 1 $\alpha(h, \varepsilon|n) \neq !$, and
- 2 $\alpha(h, \varepsilon|n+1) = b$, and
- 3 for each $m < n$, $\alpha(h, \varepsilon|m) \neq !$.



CERTAINTY IN RELIABLE INQUIRY

DEFINITION

α verifies h with certainty on ε (with respect to C) iff α produces 1 with certainty on $(h, \varepsilon) \Leftrightarrow C(\varepsilon, h)$.

DEFINITION

α refutes h with certainty on ε (with respect to C) iff α produces 0 with certainty on $(h, \varepsilon) \Leftrightarrow \neg C(\varepsilon, h)$.

DEFINITION

Decidability with certainty is simply verifiability and refutability with certainty at the same time.



EXAMPLES

- At least six bikes are broken. - Verifiable with certainty
- An even number of bikes is broken. - Verifiable in the limit

EPISTEMOLOGICAL PROPERTIES OF Q O.T. ⟨1⟩

- 1 – 1 enumeration of elements of the universe.
- Assignment of χ_A to each of them.
- Infinite sequence of 0s and 1s.
- In each step checking if finite initial segment satisfies a hypothesis (quantifier sentence).



EPISTEMOLOGICAL PROPERTIES OF Q O.T. $\langle 1 \rangle$

PROPOSITION

*Let Q be FO quantifier of type $\langle 1 \rangle$.
 Q is MON \uparrow and EXT iff it is verifiable with certainty.*

PROPOSITION

*Let Q be FO quantifier of type $\langle 1 \rangle$.
 $\neg Q$ is verifiable with certainty iff Q is falsifiable with certainty.*



EPISTEMOLOGICAL PROPERTIES OF Q O.T. $\langle 1, 1 \rangle$

PROPOSITION

*Let Q be FO CE quantifier of type $\langle 1, 1 \rangle$.
Q is persistent iff it is verifiable with certainty.*

PROPOSITION

*Let Q be FO CE quantifier of type $\langle 1, 1 \rangle$.
 \neg Q is falsifiable with certainty iff Q is verifiable with certainty.*



EXAMPLES

Determiner	verifiable	falsifiable	MON \uparrow	EXT (for $\langle 1 \rangle$)	PER (for $\langle 1, 1 \rangle$)
All	-	+	+	-	-
No	-	+	-	-	-
Some	+	-	+	+	+
At least n	+	-	+	+	+
At most n	-	+	-	+	-
Exactly n	-	-	-	+	-

TABLE: Quantifiers and their properties



PLAN

- 1 PROBLEMS
- 2 QUANTIFIERS
 - Quantifiers of type $\langle 1 \rangle$
 - Quantifiers of type $\langle 1, 1 \rangle$
- 3 COMPUTATIONAL EPISTEMOLOGY
- 4 IDENTIFIABILITY**
- 5 GENERAL QUESTION



IDENTIFIABILITY GAME

- Class of objects is chosen (e.g. class of grammars).
- Player 1 picks out one object from the class (e.g. G).
- Player 1 generates positive instances of this object, repetitions allowed (e.g. words from a language of G).
- Player 2 knows about the class, but he does not know which object is chosen.
- Player 2 has to guess which object Player 1 has in mind.




LEARNING THE SEMANTICS OF NATURAL LANGUAGE

IDENTIFIABILITY FROM TEXT IN USE

- Class of quantifiers is chosen.
- Player 1 picks one of them (Q)
- Player 2 is presented finite worlds in which Q is true.
- Player 2 has to identify Q.


NUMBER TRIANGLE REPRESENTATION

- Graphic representation of a class of CE quantifiers.
- In particular: PER.




NUMBER TRIANGLE REPRESENTATION

- Graphic representation of a class of CE quantifiers.
- In particular: PER.




NUMBER TRIANGLE REPRESENTATION

- Graphic representation of a class of CE quantifiers.
- In particular: PER.



NUMBER TRIANGLE REPRESENTATION

- Graphic representation of a class of CE quantifiers.
- In particular: PER.



TIEDE'S RESULT

THEOREM

The class of FO PER Q is identifiable from text.




PLAN

- 1 PROBLEMS
- 2 QUANTIFIERS
 - Quantifiers of type $\langle 1 \rangle$
 - Quantifiers of type $\langle 1, 1 \rangle$
- 3 COMPUTATIONAL EPISTEMOLOGY
- 4 IDENTIFIABILITY
- 5 GENERAL QUESTION




GENERAL QUESTION

RELATION BETWEEN VER/FAL HIERARCHY AND IDENTIFIABILITY




GENERAL QUESTION

RELATION BETWEEN VER/FAL HIERARCHY AND IDENTIFIABILITY




GENERAL QUESTION

RELATION BETWEEN VER/FAL HIERARCHY AND IDENTIFIABILITY




GENERAL QUESTION

RELATION BETWEEN VER/FAL HIERARCHY AND IDENTIFIABILITY




GENERAL QUESTION

RELATION BETWEEN VER/FAL HIERARCHY AND IDENTIFIABILITY




GENERAL QUESTION

RELATION BETWEEN VER/FAL HIERARCHY AND IDENTIFIABILITY




GENERAL QUESTION

RELATION BETWEEN VER/FAL HIERARCHY AND IDENTIFIABILITY




GENERAL QUESTION

RELATION BETWEEN VER/FAL HIERARCHY AND IDENTIFIABILITY




GENERAL QUESTION

RELATION BETWEEN VER/FAL HIERARCHY AND IDENTIFIABILITY




GENERAL QUESTION

RELATION BETWEEN VER/FAL HIERARCHY AND IDENTIFIABILITY



GENERAL QUESTION

RELATION BETWEEN VER/FAL HIERARCHY AND IDENTIFIABILITY



CONCLUSIONS AND FUTURE WORK

- Epistemological role of monotonicity - additional explanation.
- Verification less difficult than falsification?
- Check connections between persistence and comprehension.
- Investigate relationship between identifiability and decidability: learning of NL semantics; new conditions of identifiability.



THANK YOU!